

BEST AVAILABLE COPY

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
15 January 2004 (15.01.2004)

PCT

(10) International Publication Number
WO 2004/004697 A2

(51) International Patent Classification⁷: **A61K 31/00** (74) Agents: **RAMBELLI, Paolo et al.; Jacobacci & Partners SpA, Corso Regio Parco 27, I-10152 Torino (IT).**

(21) International Application Number:
PCT/IB2003/002632

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(22) International Filing Date: 3 July 2003 (03.07.2003)

(25) Filing Language: Italian

(26) Publication Language: English

(30) Priority Data:
TO2002A000582 4 July 2002 (04.07.2002) IT

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BE, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicants (*for all designated States except US*): **UNIVERSITA' DEGLI STUDI DI FERRARA [IT/IT]; Via Savonarola, 9, I-44100 Ferrara (IT). ASSOCIAZIONE VENETA PER LA LOTTA ALLA [IT/IT]; c/o Centro Microcitemia Dell'Azienda ULSS 18, I-45100 Rovigo (IT).**

Declaration under Rule 4.17:

— *of inventorship (Rule 4.17(iv)) for US only*

Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 2004/004697 A2

(54) Title: A NOVEL USE OF RAPAMYCIN AND STRUCTURAL ANALOGUES THEREOF

(57) Abstract: The use of rapamycin and its structural analogues for the preparation of a medicament for the therapeutic treatment of beta-thalassaemia is described.

A novel use of rapamycin and structural analogues thereof

The present invention relates to the use of rapamycin and its structural analogues for the therapeutic treatment of beta-thalassaemia.

The existence of substances capable of inducing the expression of the gene for gamma-globin and the biosynthesis of foetal haemoglobin (HbF) in adult subjects has been known for some time (1). In the majority of cases, those substances are also capable of activating or potentiating the transcription of genes for embryonal and foetal globins in experimental model systems.


Recently, for example, numerous DNA-binding molecules have been described that have the capacity to bring about an increase in the expression of genes for gamma-globin (2). Among these there may be mentioned cisplatin and analogues of cisplatin, mithramycin and chromomycin, tallimustine (3). Those molecules are efficient modulators of the expression of genes for gamma-globin.

In human beings, activation of the transcription of genes for gamma-globins in adult subjects leads to the production of foetal haemoglobin mimicking the phenotype HPFH (High Persistence of Foetal Haemoglobin) which confers a favourable clinical picture on patients suffering from beta-thalassaemia also in homozygotic form (4). A therapy providing for the use of those molecules in the treatment of patients suffering from beta-thalassaemia could therefore make those subjects less dependent on transfusion therapy (5).

The present invention is based on the need for novel modifiers of the transcription process that can be used in the treatment of beta-thalassaemia and that have a high level of

induction of the expression of gamma-globin genes and a low level of cytotoxicity.

We have surprisingly found that rapamycin, an antibiotic which is described in the prior art and the structural formula of which is illustrated hereinafter, satisfies those requirements.



RAPAMYCIN

In particular, it has been found that rapamycin and its structural analogues are capable of activating the expression of the gene for human gamma-globin.

Rapamycin is an antibiotic produced by *Streptomyces hygroscopicus* and originally described as an anti-fungal agent (6).

The chemical synthesis of rapamycin and its structural analogues - for example 7-(N-hydroxy)-acyl, -carbamoyl and -ureide analogues - has been described by various research groups (see, for example, the bibliographical references 7 and 8).

Rapamycin is currently the active ingredient of drugs (Sirolimus, Rapamune, antibiotic AY22989) used as anti-inflammatory agents (9, 10) in the treatment of atopical dermatitis (9), allergic contact dermatitis (9) and psoriasis (10), as anti-tumour agents (11, 12) and anti-fungal agents (6). In addition, rapamycin is widely used in therapy as a drug capable of retarding the rejection process (13) in heart (14), kidney (15) and liver (16) transplants.

However, the capacity of rapamycin and its structural analogues to activate the expression of the human gamma-globin gene has not been described hitherto, nor this activity is attributable to its known effects indicated above.

A first subject of the present invention is therefore the use of rapamycin and its structural analogues for the preparation of a medicament for the therapeutic treatment of beta-thalassaemia.

As has been recently described (17, 18), a combined treatment with various modifiers of the transcription process would also permit a further increase in the expression of genes for gamma-globin.

Therefore, a second subject of the present invention is the use of rapamycin or its structural analogues in combination with at least one further modifier of the transcription process for the preparation of a medicament for the treatment of beta-thalassaemia.

According to a preferred embodiment, said further modifier of the transcription process is selected from the group consisting of cytosine arabinoside, retinoic acid, plicamycin, hydroxyurea, guanine, guanosine triphosphate (GTP), guanosine diphosphate (GDP) and guanosine monophosphate (GMP); of

these, cytosine arabinoside and retinoic acid are more preferred.

The activity of rapamycin as an inducer of erythroid cell differentiation and the production of mRNA for gamma-globin was evaluated in human cell cultures. The results of this study are illustrated in the Tables given in the section relating to the Examples. The data obtained indicate that the activity of rapamycin is greater than that of reference drugs (for example, hydroxyurea in the induction of HbF); it has also been ascertained that the cytotoxic effect which may be encountered is much lower than that of hydroxyurea.

The Examples which follow are provided for the purposes of illustration and are not intended to limit the scope of the invention in any way.

Example 1

The biological activity of rapamycin was evaluated by examining the capacity of that compound to modulate the expression of genes for gamma-globin in the human cell line K562, which is capable of undergoing erythroid differentiation by expressing genes for gamma-globin if subjected to treatment with modifiers of the biological response that are suitable for the purpose (3, 17, 18). The level of differentiation was evaluated by analysing the positive reaction of the cells to benzidine (3). Some of the data obtained are given in Table 1, which shows the results of six independent experiments (mean \pm SD). As will be readily appreciated, rapamycin is capable of bringing about an increase in the percentage of K562 cells that react positively to benzidine (40-45% in the treated cells, compared with 4.1% of the control K562 cells). The production of haemoglobin was evaluated by electrophoresis on cellulose acetate and by staining the gel with a solu-

tion based on benzidine/H₂O₂ (3). The chief haemoglobin produced by K562 cells treated with rapamycin was Hb Portland (alpha2gamma2). The expression of genes coding for gamma-globins was evaluated by quantitative RT-PCR (reverse transcriptase PCR) (3). The data obtained demonstrate an increase in the intracytoplasmic accumulation of mRNA for gamma-globin. These evaluations were carried out after 6 days of induction with 20 nM rapamycin. The results of the quantitative RT-PCR are shown in Table 2.

Table 1

Compound	Concentration (nM)	^a Erythroid differentiation (%)
-	-	4.1±3.7
Rapamycin	20	45.5±5.8

^a Erythroid differentiation = percentage of K562 cells that react positively to benzidine (mean ± SD of 6 experiments).

Table 2

Compound	Concentration (nM)	^b mRNA for gamma-globin
-	-	1
Rapamycin	20	2.4

^b The accumulation of RNA for gamma-globin is given in the Table as an increase compared with that of untreated control K562 cells. The technique used was that of quantitative RT-PCR (21, 22) using the following primer and probe oligonu-

cleotides: gamma-globin forward primer, 5'-TGG CAA GAA GGT GCT GAC TTC-3'; gamma-globin reverse primer, 5'-TCA CTC AGC TGG GCA AAG G-3'; gamma-globin probe 5'-FAM-TGG GAG ATG CCA TAA AGC ACC TGG-TAMRA-3' (FAM = 6-carboxy fluorescein, TAMRA = 6-carboxy-N,N,N',N'-tetramethylrhodamine).

Example 2

In order to check whether rapamycin was capable, in addition to inducing K562 cell differentiation, of stimulating the production of mRNA for gamma-globin in human erythroid precursors isolated from peripheral blood, the technique published by Fibach et al. (19, 20) was used. This technique provides for two stages: in the first stage, the cells isolated from peripheral blood of a subject who is healthy or suffering from a haemopoietic pathology, such as sickle cell anaemia or beta-thalassaemia, are sown in a culture medium to which 10% of conditioned medium derived from the vesicle carcinoma cell line 5637 has been added. The second stage consists in cultivating the isolated cells in a suitable culture medium, supplemented by the erythropoietin hormone, 30% foetal bovine serum, 2-mercaptoethanol, albumin, glutamine and desamethasone in order to permit the proliferation and maturing of stem cells of the erythroid type. In this stage the cells can be treated with potential HbF inducers.

For example, in this system it was demonstrated that hydroxyurea, an inhibitor of DNA synthesis currently used in the experimental therapy of beta-thalassaemia, is capable of bringing about the production of HbF.

The results obtained with rapamycin demonstrated an increase in the production of mRNA for gamma-globin in cells treated with rapamycin compared with the same untreated cells (3.75

times), which increase is greater than that which may be encountered using hydroxyurea (Table 3).

Table 3

Compound	Concentration (μ M)	^a mRNA for gamma-globin
-	-	1
Rapamycin	3	3.73
Hydroxyurea	120	2.25

^a The accumulation of RNA for gamma-globin is given in the Table as an increase compared with that of untreated control erythroid precursors. The technique used was that of quantitative RT-PCR (21, 22) using the primer and probe oligonucleotides described in Table 2.

Bibliographical references

1. Rodgers GP, Rachmilewitz EA, British J. Haematology, 91:263-268, 1995.
2. Rochette J, Craig JE and Thein SL, Blood Reviews, 8: 213-224, 1994.
3. Bianchi N, Chiarabelli C, Borgatti M, Mischiati C, Fibach E, Gambari R. Br J Haematol. 113(4):951-61, 2001.
4. Dover, G.J., Brusilow, S and Samid D, New England Journal of Medicine, 327: 569-570, 1992.

5. Ikuta, T., Atweh, G., Boosalis, V., White, G.L., De Fonseca, S., Boosalis, M., Faller, D.V., Perrine, S.P., Annals of New York Academy of Sciences, 850:87-99, 1998.

6. Kahan BD. Sirolimus: a comprehensive review. Expert Opin Pharmacother. 2001 Nov;2(11):1903-17.

7. Dickman DA, Ding H, Li Q, Nilius AM, Balli DJ, Ballaron SJ, Trevillyan JM, Smith ML, Seif LS, Kim K, Sarthy A, Goldman RC, Plattner JJ, Bennani YL. Bioorg Med Chem Lett. 10(13):1405-8, 2000.

8. Dell CP. Curr Med Chem. 5(3):179-94, 1998.

9. Trautmann A, Akdis M, Schmid-Grendelmeier P, Disch R, Brocker EB, Blaser K, Akdis CA. J Allergy Clin Immunol. 108(5):839-46, 2001.

10. Reitamo S, Spuls P, Sassolas B, Lahfa M, Claudy A, Griffiths CE. Br J Dermatol. 145(3):438-45, 2001.

11. Guba M, von Breitenbuch P, Steinbauer M, Koehl G, Flegel S, Hornung M, Bruns CJ, Zuelke C, Farkas S, Anthuber M, Jauch KW, Geissler EK. Nat Med. 8(2):128-35, 2002.

12. Garber K. J Natl Cancer Inst. 93(20):1517-9, 2001.

13. Benito AI, Furlong T, Martin PJ, Anasetti C, Appelbaum FR, Doney K, Nash RA, Papayannopoulou T, Storb R, Sullivan KM, Witherspoon R, Deeg HJ. Transplantation. 72(12):1924-9, 2001.

14. Radovancevic B, El-Sabrout R, Thomas C, Radovancevic R, Frazier OH, Van Buren C. 33(7-8):3221-2, 2001.

15. Podbielski J, Schoenberg L. Prog Transplant. 11(1):29-32, 2001.

16. Nishida S, Pinna A, Verzaro R, Levi D, Kato T, Khan F, Nery J, Weppler D, Tzakis A. Transplant Proc. 33(1-2):1495, 2001.

17. Bianchi N, Ongaro F, Chiarabelli C, Gualandi L, Mischiati C, Bergamini P, Gambari R. Biochem Pharmacol. 60:31-40, 2000.

18. Bianchi N, Osti F, Rutigliano C, Ginanni Corradini F, Borsetti E, Tomassetti M, Mischiati C, Feriotto G e Gambari R, British Journal of Haematology, 104:258-263, 1999.

19. Fibach E. Hemoglobin, 22: 445-458, 1998.

20. Fibach E, Burke KP, Schechter AN, Noguchi CT & Rodgers GP. Blood, 81: 1630-1635, 1993.

21. Heid CA, Stevens J, Livak KJ & Williams PM. Genome Research, 6: 986-994, 1996.

22. Gibson UE, Heid CA & Williams PM. Genome Research, 6: 995-1001, 1996.

CLAIMS

1. Use of rapamycin and its structural analogues for the preparation of a medicament for the therapeutic treatment of beta-thalassaemia.
2. Use according to claim 1, wherein the structural analogue is selected from the group comprising 7-(N-hydroxy)-acyl analogues, carbamoyl analogues and ureide analogues of rapamycin.
3. Use according to claim 1 or 2, wherein the rapamycin or the structural analogue is in combination with at least one further modifier of the transcription process selected from the group consisting of cytosine arabinoside, retinoic acid, plicamycin, mithramycin, hydroxyurea, guanine, guanosine triphosphate (GTP), guanosine diphosphate (GDP) and guanosine monophosphate (GMP).

THIS PAGE BLANK (USPTO)

(19) World Intellectual Property Organization International Bureau



(43) International Publication Date
15 January 2004 (15.01.2004)

PCT

(10) International Publication Number
WO 2004/004697 A3

(51) International Patent Classification⁷: **A61K 31/453**, 31/203, 31/7068, 31/704, 31/17, 31/708

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(21) International Application Number:
PCT/IB2003/002632

(22) International Filing Date: 3 July 2003 (03.07.2003)

(25) Filing Language: Italian

(26) Publication Language: English

(30) Priority Data:
TO2002A000582 4 July 2002 (04.07.2002) IT

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicants (*for all designated States except US*): UNIVERSITA' DEGLI STUDI DI FERRARA [IT/IT]; Via Savonarola, 9, I-44100 Ferrara (IT). ASSOCIAZIONE VENETA PER LA LOTTA ALLA [IT/IT]; c/o Centro Microcitemia Dell'Azienda ULSS 18, I-45100 Rovigo (IT).

Declaration under Rule 4.17:

— *of inventorship (Rule 4.17(iv)) for US only*

Published:

— *with international search report*
— *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*

(88) Date of publication of the international search report:
25 March 2004

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): BIANCHI, Nicoletta [IT/IT]; Via G. Matteotti 10/A, I-44020 Mezzogoro (IT). BORGATTI, Monica [IT/IT]; Via Menotti Cova 6, I-44100 Ferrara (IT). GAMBARI, Roberto [IT/IT]; Via Normandia, 17, I-40132 Bologna (IT). MISCHIATI, Carlo [IT/IT]; Via Don Minzoni, 99, I-45030 Rovigo (IT).

(74) Agents: RAMBELLI, Paolo et al.; Jacobacci & Partners SpA, Corso Regio Parco 27, I-10152 Torino (IT).

WO 2004/004697 A3

(54) Title: A NOVEL USE OF RAPAMYCIN AND STRUCTURAL ANALOGUES THEREOF

(57) Abstract: The use of rapamycin and its structural analogues for the preparation of a medicament for the therapeutic treatment of beta-thalassaemia is described.

INTERNATIONAL SEARCH REPORT

Inte... nal Application No
PCT/IB 03/02632

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/453 A61K31/203 A61K31/7068 A61K31/704 A61K31/17
A61K31/708

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

MEDLINE, BIOSIS, EPO-Internal, WPI Data, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JOHNSTON JULIE ET AL: "Treatment of beta-thalassemia in the mouse by regulated expression of AAV-encoded erythropoietin" BLOOD, vol. 98, no. 11 Part 2, 16 November 2001 (2001-11-16), page 410b XP009023880 43rd Annual Meeting of the American Society of Hematology, Part 2; Orlando, Florida, USA; December 07-11, 2001 ISSN: 0006-4971	1-3
Y	the whole document --- -/-	3

Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

15 January 2004

Date of mailing of the international search report

28/01/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel: (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Loher, F

INTERNATIONAL SEARCH REPORT

International Application No
PCT/IB 03/02632

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 01 68147 A (BIANCHI NICOLETTA ;FERIOTTO GIORDANA (IT); GAMBARI ROBERTO (IT); M) 20 September 2001 (2001-09-20) page 1, paragraph 1 – paragraph 3 page 5, paragraph 1 -----	3

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 03/02632

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
WO 0168147	A 20-09-2001	IT	T020000234 A1	13-09-2001
		AU	5827401 A	24-09-2001
		BR	0109365 A	24-12-2002
		WO	0168147 A2	20-09-2001
		EP	1263475 A2	11-12-2002
		US	2003073660 A1	17-04-2003

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- BLACK BORDERS**
- IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- FADED TEXT OR DRAWING**
- BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- SKEWED/SLANTED IMAGES**
- COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- GRAY SCALE DOCUMENTS**
- LINES OR MARKS ON ORIGINAL DOCUMENT**
- REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)